

Climate Forcing and Feedback from the Terrestrial Carbon Cycle and Land Cover Change

Gordon Bonan
National Center for Atmospheric Research
Boulder, Colorado

19 January 2010
Department of Environmental Sciences
University of Virginia
Charlottesville, Virginia



Evolution of climate science

Atmospheric sciences

Atmospheric general circulation models

- Atmospheric physics and dynamics
- Prescribed sea-surface temperature and sea ice
- Bulk formulation of surface fluxes without vegetation
- Bucket model of soil hydrology (1970s)

Land surface models

- Surface energy balance
- Hydrologic cycle
- Plant canopy (1980s)

Oceanography

Ocean general circulation models

- Physics, dynamics

Atmospheric & oceanic sciences

Global climate models

- Atmosphere
- Land and vegetation
- Ocean
- Sea ice (1990s)

Ocean ecosystem models

- Biogeochemical cycles

Ecology


Terrestrial ecosystem models

- Biogeochemical cycles (C-N-P)
- Vegetation dynamics
- Wildfire
- Land use (1990s)

Earth system science

Earth system models

- Physics, chemistry, biology
- Humans, socioeconomics (2000s)



1. Introduction

Climate of the 21st century


Multi-model mean surface warming (relative to 1980-1999) for the scenarios A2, A1B and B1

Multi-model mean warming and uncertainty for 2090 to 2099 relative to 1980 to 1999:

A2: +3.4°C (2.0°C to 5.4°C)

A1B: +2.8°C (1.7°C to 4.4°C)


B1: +1.8°C (1.1°C to 2.9°C)




Meehl et al. (2007) in *Climate Change 2007: The Physical Science Basis*, Solomon et al., Eds., 747-845

For 5th assessment report

- o Land cover change and the carbon cycle as climate forcings and feedbacks
- o Can ecosystems be managed to mitigate climate change?



Multiple competing influences of ecosystems





Boreal forest - menace to society - no need to promote conservation



Temperate forest - reforestation and afforestation?



Tropical rainforest - planetary savior - promote avoided deforestation, reforestation, or afforestation



Biofuel plantations to lower albedo and reduce atmospheric CO_2




Outline of talk

1. Introduction
2. Representing ecosystems in climate models
3. Carbon cycle and climate
 - Concentration-carbon feedback (CO_2 fertilization)
 - Climate-carbon feedback (temperature)
 - Nitrogen cycle
4. Land use and land cover change
 - 4a. Biogeochemical*
Land use carbon flux
 - 4b. Biogeophysical*
Albedo and evapotranspiration
5. Climate change mitigation

2. Models

The Earth system



(IPCC 2007)

Climate models use mathematical formulas to simulate the **physical**, **chemical**, and **biological** processes that drive Earth's climate

A typical climate model consists of coupled models of the **atmosphere**, **ocean**, **sea ice**, and **land**

Land is represented by its **ecosystems**, **watersheds**, **people**, and **socioeconomic drivers** of environmental change

The model provides a comprehensive understanding of the processes by which people and ecosystems affect, adapt to, and mitigate global change

2. Models

The Community Land Model

Fluxes of energy, water, and carbon and the dynamical processes that alter these fluxes

Oleson et al. (2004) NCAR/TN-461+STR

Oleson et al. (2008) JGR, 113,
doi:10.1029/2007JG000563


Stöckli et al. (2008) JGR, 113,
doi:10.1029/2007JG000562

Spatial scale

1.25° longitude × 0.9375° latitude
(288 × 192 grid)


Temporal scale

- o 30-minute coupling with atmosphere
- o Seasonal-to-interannual (phenology)
- o Decadal-to-century climate (disturbance, land use, succession)
- o Paleoclimate (biogeography)



Land surface heterogeneity

Subgrid land cover and plant functional types




CLM represents a model grid cell as a mosaic of up to 6 primary land cover types. Vegetated land is further represented as a mosaic of plant functional types



Local land use is spatially heterogeneous




Patchwork of agricultural land, Colorado (NCAR)



Settlement and deforestation surrounding Rio Branco, Brazil (10°S, 68°W) in the Brazilian state of Acre, near the border with Bolivia. The large image covers an area of 333 km x 333 km (NASA/GSFC/LaRC/JPL)





Global land use is abstracted to the fractional area of crops and pasture



Foley et al. (2005) Science 309:570-574

Flux tower measurements - temperate deciduous forest

Morgan Monroe State Forest,
Indiana




CLM3.0 - dry soil, low latent heat flux, high sensible heat flux
 CLM3.5 - wetter soil and higher latent heat flux

Annual net primary production

Ecosystem Model-Data Intercomparison
(EMDI) compilation of observations

- Class A (81 sites)
- Class B (933 sites)

NPP extracted for each model grid cell
corresponding to a measurement location



2. Models

Integrate ecological studies with earth system models

Environmental Monitoring



Eddy covariance flux tower
(courtesy Dennis Baldocchi)

Experimental Manipulation



Soil warming, Harvard Forest




CO₂ enrichment, Duke Forest



Hubbard Brook
Ecosystem Study

Test model-generated hypotheses of earth system functioning with observations


Modeling the Climate System



Planetary energetics
Planetary ecology
Planetary metabolism



Effect of climate change on carbon cycle



Friedlingstein et al. (2006) J Climate 19:3337-3353

Climate-carbon cycle feedback

11 carbon cycle-climate models of varying complexity

All models have a positive climate-carbon cycle feedback (20 ppm to >200 ppm)

Atmospheric carbon increases compared with no climate-carbon cycle feedback, while land carbon storage decreases


Prevailing model paradigm

CO_2 fertilization enhances carbon uptake, diminished by decreased productivity and increased soil carbon loss with warming

But what about the nitrogen cycle and land use?

Prevailing modeling paradigm

CO_2 fertilization enhances carbon uptake, diminished by decreased productivity and increased soil carbon loss with warming



$$\Delta C_L = \beta_L \Delta C_A$$

$$\Delta C_L = \beta_L \Delta C_A + \gamma_L \Delta T$$

$\beta_L > 0$: concentration-carbon feedback (Pg C ppm^{-1})


$\gamma_L < 0$: climate-carbon feedback (Pg C K^{-1})

Carbon-nitrogen interactions


Reduces concentration-carbon feedback (β_L)
Changes sign of climate-carbon feedback (γ_L)

Sokolov et al. (2008) *J Climate* 21:3776-3796
Thornton et al. (2009) *Biogeosci* 6:2099-2120

- Nitrogen limitation reduces the CO_2 fertilization gain in productivity
- Greater N mineralization with warming stimulates plant growth




Land biosphere response to CO_2



Land biosphere response to temperature

Thick solid line is with preindustrial nitrogen deposition
Thick dashed line is with anthropogenic nitrogen deposition
Thin gray lines are C4MIP models

Multi-scale carbon cycle evaluation



Randerson et al. (2009) *GCB* 15:2462-2484

1. Test model-generated hypotheses of earth system functioning with observations
2. Model experimentation to inform key research needs

Quantifying carbon-nitrogen feedbacks in CLM4

Annual Mean Forcings (Land Only) for Control and Experiment Simulations

				Land Use	
Simulations	Atmos. CO ₂ [ppm]	Temperature [K]	N deposition [Tg N yr ⁻¹]	Cropland [10 ⁶ km ²]	Wood harvest [10 ⁶ km ² yr ⁻¹]
Control	328.6	280.8	48.5	14.0	0
Experiments					
1973-77	331.0	280.9	51.2	14.1	0.14
2000-04	372.8	281.8	63.9	15.2	0.22
Change	41.8	0.9	12.7	1.1	0.08

Forcings are constant for control simulations and vary with time for experiment simulations. Shown are the 1973-1977 and 2000-2004 means and the temporal change.

Quantifying carbon-nitrogen feedbacks in CLM4


Carbon fluxes 1973 - 2004

	C	CN _{ndep}	GCP
Land use (Pg C yr ⁻¹)	1.8	1.8	1.5
Land sink (Pg C yr ⁻¹)	2.5	1.8	2.0 - 2.4

Global Carbon Project (www.globalcarbonproject.org)

Le Quéré et al. (2009) *Nature Geosci* 2:831-836

Time series of annual land uptake



Quantifying carbon-nitrogen feedbacks in CLM4

β_L and γ_L Calculated for Carbon-Only and Carbon-Nitrogen Simulations

	Without HLCC		With HLCC
β_L (Pg C ppm ⁻¹)	Constant Climate	Climate Change	Climate Change
C	0.94	0.94	0.92
CN _{ndep}	0.25	0.26	0.25
γ_L (Pg C K ⁻¹)	Constant CO ₂	Increasing CO ₂	Increasing CO ₂
C	-11.7	-11.7	-11.0
CN _{ndep}	-0.9	-0.2	0.2

C mean β_L is 3.7 times greater than CN_{ndep} mean (i.e., 73% reduction in β_L)

CN_{ndep} reduces carbon loss with climate change, i.e., γ_L increases

Quantifying carbon-nitrogen feedbacks in CLM4

Carbon budget analysis (Pg C yr^{-1})

$$\Delta C_L' = \Delta C_L^{\text{HIST}} + \Delta \Delta C_L^{\text{CONC}} + \Delta \Delta C_L^{\text{CLIM}} + \Delta \Delta C_L^{\text{NDEP}} + \Delta \Delta C_L^{\text{HLCC}}$$

Simulation	ΔC_L	$\Delta C_L'$	ΔC_L^{HIST}	$\Delta \Delta C_L$			
				CONC	CLIM	NDEP	HLCC
C	0.62	0.62	1.54	1.43	-0.37	0.00	-1.97
CN _{ndep}	-0.13	-0.11	1.22	0.38	0.01	0.19	-1.92
CN _{ndep} - C	-0.75	-0.73	-0.32	-1.04	0.38	0.19	0.05

C: CONC feedback is four times greater than CLIM feedback

CN_{ndep}: decrease in CONC uptake is three times greater than reduction in CLIM loss

The influence of nitrogen on the concentration-carbon feedback is of greater importance for near-term climate change simulations than its effect on the climate-carbon feedback

The land use carbon flux greatly exceeds these carbon-nitrogen biogeochemical feedbacks


Representing land use and land cover change

1. For IPCC AR5 land use and land cover change are to be described consistently with Representative Concentration Pathways (RCP) scenarios
2. All pathways share the same historical trajectory to 2005. After 2005 they diverge following own representative pathway.
3. For the historical period and for each RCP, land use that results in land cover change is described through annual changes in four basic land units:
 - Primary Vegetation (V)
 - Secondary Vegetation (S)
 - Cropping (C)
 - Pasture (P)
4. Harvesting of biomass is also prescribed for both primary and secondary vegetation land units
5. George Hurtt and colleagues at University of New Hampshire are harmonizing the historical and RCP data (luh.unh.edu)


4. Land use

Historical land cover change, 1850 to 2005


Tree PFTs




Crop PFT



Shrub PFTs



Grass PFTs




(datasets by Lawrence & Feddema)

4. Land use

Future land cover change, 2005 to 2100

MESSAGE (RCP 8.5 $W m^{-2}$)



MINICAM (RCP 4.5 $W m^{-2}$)





IMAGE (RCP 2.6 $W m^{-2}$)




AIM (RCP 6.0 $W m^{-2}$)

(In development)

4. Land use

Future land cover change, 2005 to 2100 (RCPs)

MESSAGE (RCP 8.5 $W m^{-2}$)



MINICAM (RCP 4.5 $W m^{-2}$)






IMAGE (RCP 2.6 $W m^{-2}$)



AIM (RCP 6.0 $W m^{-2}$)


(In development)

Land use - wood harvest



(datasets by Lawrence & Feddema)


Carbon flux to wood products



Wood harvesting

Land cover change
(e.g., deforestation)


(simulations by Sam Levis)



Shevliakova et al. (2009) *GBC*, 23, GB2022, doi:10.1029/2007GB003176


4. Land use

Land use carbon flux to atmosphere



Wood harvesting


Land cover change
(e.g., deforestation)




The Woods Hole Research Center

4. Land use

Land use carbon flux to atmosphere



The LUCID intercomparison study



Multi-model ensemble of global land use climate forcing (1992-1870)

Seven climate models of varying complexity with imposed land cover change (1992-1870)

Pitman et al. (2009) *GRL*, 36, L14814, doi:10.1029/2009GL039076

Models

Atmosphere - CAM3.5

Land - CLM3.5 + new datasets for present-day vegetation + grass optical properties

Ocean - Prescribed SSTs and sea ice

Experiments

30-year simulations ($CO_2 = 375$ ppm, SSTs = 1972-2001)

PD - 1992 vegetation

PDv - 1870 vegetation

30-year simulations ($CO_2 = 280$ ppm, SSTs = 1871-1900)

PI - 1870 vegetation

PIv - 1992 vegetation


No irrigation

5-member ensembles each

Total of 20 simulations and 600 model years

4. Land use


The LUCID intercomparison study




4. Land use

The LUCID intercomparison study


Latent Heat Flux Difference
IPSL-ORCHIDEE




ARPEGE-ISBA




CCSM-CLM




CCAM-CABLE




ECEarth



SPEEDY-LPJ



ECHAM5-JSBACH



Change in JJA latent heat flux (W m^{-2}) resulting from land cover change (PD - PDv)

4. Land use



Albedo forcing, 1992-1870

Present Day - 1870 DJF Surface Albedo

(-)

Present Day - 1870 MAM Surface Albedo

(-)







Present Day - 1870 JJA Surface Albedo

(-)

Present Day - 1870 SON Surface Albedo

(-)




4. Land use

Near-surface temperature, 1992-1870


Present Day - 1870 DJF Atmospheric Temperature

(°C)




Present Day - 1870 MAM Atmospheric Temperature

(°C)






Present Day - 1870 JJA Atmospheric Temperature

(°C)





Present Day - 1870 SON Atmospheric Temperature

(°C)



4. Land use


Atmospheric feedbacks



Climate models simulate the large-scale response and include feedbacks with the atmosphere:

- o Increased rainfall enhances latent heat flux
- o Increased cloudiness reduces solar radiation
- o Reduced PBL height

Flux towers measure local response



4. Land use




Land cover change offsets greenhouse gas warming

Land cover change with $CO_2 = 375$ ppm (1992)

Land cover change with $CO_2 = 280$ ppm (1870)



Present Day - 1870 JJA Atmospheric Temperature ($^{\circ}C$)

1992 - 1870 JJA Atmospheric Temperature ($^{\circ}C$)



CO_2 forcing with 1870 land cover

1992 - 1870 JJA Atmospheric Temperature ($^{\circ}C$)




Cropland increases surface albedo

Monthly shortwave surface albedo for dominant US land cover types in the Northeast (b) and Southeast (d)

Jackson et al. (2008) Environ Res Lett, 3, 044006 (doi:10.1088/1748-9326/3/4/044006)

Forest masking

Cropland has a high winter and summer albedo compared with forest




Land cover change and evapotranspiration

Prevailing model paradigm

Crops

Low latent heat flux because of:


- Low roughness
- Shallow roots decrease soil water availability



Trees

High latent heat flux because of:


- High roughness
- Deep roots allow increased soil water availability



Tropical forest - cooling from higher surface albedo of cropland and pastureland is offset by warming associated with reduced evapotranspiration

Temperate forest - higher albedo leads to cooling, but changes in evapotranspiration can either enhance or mitigate this cooling

Reforestation cools climate



Annual mean temperature change

	OF to PP	OF to HW
Albedo	+0.9°C	+0.7°C
Ecophysiology and aerodynamics	-2.9°C	-2.1°C


Forest

Lower albedo (+)

Greater leaf area index,
aerodynamic conductance, and
latent heat flux (-)

Can Ameriflux provide insights?

NCEAS "Forest and Climate Policy"
working group



Crops

Mead irrigated sites have highest LH

LH varies with crop rotation

LH varies with crop type (winter wheat)


Ecosystems

- Reforestation, afforestation, avoided deforestation
- Biofuels
- Biogeophysics and biogeochemistry (albedo, ET, carbon)

Urban planning and design

- White roofs
- Greenspaces

Average summer difference in the urban minus rural air temperature with roof albedos maximized





Oleson et al. (2010) GRL, in press

5. Mitigation

Land use choices affect 21st century climate


Future IPCC SRES land cover scenarios for NCAR LSM/PCM

a) Present day land cover




A2 - Widespread agricultural expansion with most land suitable for agriculture used for farming by 2100 to support a large global population

b) B1 2050 land cover




d) A2 2050 land cover





B1 - Loss of farmland and net reforestation due to declining global population and farm abandonment in the latter part of the century

c) B1 2100 land cover



e) A2 2100 and cover





Land use choices affect 21st century climate




SRES B1 JJA reference height temperature

SRES A2

2050



2100



Change in temperature due to land cover

B1

- Weak temperate warming
- Weak tropical warming

A2

- Temperate cooling
- Tropical warming

Conclusions

The ecology of climate models

- Detailed representation of ecosystems
- Allows exploration of ecological feedbacks and mitigation options

Carbon cycle

- CO_2 fertilization enhances carbon gain, diminished by carbon loss with warming
- N cycle reduces the concentration-carbon gain and decreases climate-carbon loss
- The CO_2 fertilization effect is larger than the climate feedback effect

Land use and land cover change

Biogeochemistry

- Wood harvest flux is important
- Uncertainty in land use flux may be greater than the N-cycle feedback

Biogeophysics

- Higher albedo of croplands cools climate
- Less certainty about role of latent heat flux
- Implementation of land cover change (spatial extent, crop parameterization) matters